

Introduction to Gauge/Gravity Duality

Examples II

To hand in Thursday 4th November in the examples class

I. Coordinates of AdS_{d+1} .

Lorentzian AdS_{d+1} can be defined by the locus

$$-L^2 = \eta_{ab}X^aX^b = -\left(X^{d+1}\right)^2 - \left(X^0\right)^2 + \sum_{i=1}^d \left(X^i\right)^2, \quad (1)$$

where $X \in \mathbb{R}^{2,d}$ and $ds^2 = \eta_{ab}X^aX^b$ with $\eta = \text{diag}(-1, 1, 1, \dots, 1, -1)$. In the following we parametrize the locus (1) in different ways.

a) Draw a picture of AdS_2 embedded in $\mathbb{R}^{2,1}$. (2 points)

b) The *global coordinates* (ρ, τ, Ω_i) are defined by

$$\begin{aligned} X^{d+1} &= L \cosh \rho \sin \tau, \\ X^0 &= L \cosh \rho \cos \tau, \\ X^i &= L \sinh \rho \Omega_i, \end{aligned}$$

with $i = 1, \dots, d$ and $\sum_{i=1}^d \Omega_i^2 = 1$. Using this parametrization calculate the induced metric $ds^2 = g_{\mu\nu}dx^\mu dx^\nu$ (where $x^\mu \in \{\rho, \tau, \Omega_i\}$) for AdS_{d+1} in global coordinates. (3 points)

c) Replace ρ by $r \equiv L \sinh \rho$ and show that the metric can be written in the form

$$ds^2 = -H(r)dt^2 + H(r)^{-1}dr^2 + r^2 d\Omega_{d-1}^2,$$

where $d\Omega_{d-1}^2 = \sum_{i=1}^d d\Omega_i d\Omega_i$ is the metric of the unit $(d-1)$ -sphere, S^{d-1} . (2 points)

d) The *Poincare patch coordinates* (x^i, u) with $i = 1, \dots, d$ are defined by

$$\begin{aligned} X^{d+1} + X^d &= u, \\ -X^{d+1} + X^d &= v, \\ X^i &= \frac{u}{L}x^i. \end{aligned}$$

Use the defining equation (1) to eliminate v in terms of u and x^i and show that the induced metric for (u, x^i) with $i = 1, \dots, d$ takes the form

$$ds^2 = L^2 \frac{du^2}{u^2} + \frac{u^2}{L^2} dx^i dx_i.$$

Finally introduce $z = \frac{L^2}{u}$ and show that the metric is given by

$$ds^2 = \frac{L^2}{z^2} (dz^2 + dx^i dx_i)$$

Which part of the AdS spacetime is not covered by these coordinates (Hint: z takes only positive values (Why?)). (3 points)