

From Here to Infinity: Extending Frontiers in Cosmology
A Weekend Symposium to Celebrate the Sixtieth Birthday of Joseph Silk
Oxford, England, December 13-15, 2002

Massive Neutrinos

News from the Ghost Particles



Georg Raffelt
Max-Planck-Institut für Physik, München, Germany

Status of Evidence for Neutrino Oscillations

System	Atmospheric	Solar	LSND
Channel	$\nu_\mu \rightarrow \nu_\tau$	$\nu_e \rightarrow \nu_{\mu\tau}$	$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$
$\delta m^2 / \text{eV}^2$	$(1.5 - 4) \times 10^{-3}$	LMA $(0.2 - 2) \times 10^{-4}$	0.2-2 or 6.5
$\sin^2 2\theta$	0.9-1	0.2-0.6	0.001-0.03
Status	Established	Established	Unconfirmed
Test	Long Baseline	KamLAND 12/2002	MiniBooNE 2004 ?
Implication	Mutually inconsistent, even with a sterile neutrino Evidence for physics beyond flavor oscillations (CPT violation ...) ?		
Simplest interpretation	Three mass eigenstates with $m_1 \ll m_2 \ll m_3 \sim 50 \text{ meV}$ (hierarchical) $m_1 \sim m_2 \sim m_3 \gg 50 \text{ meV}$ (degenerate)		Experimental or Statistical Fluke

Status of Evidence for Neutrino Oscillations

**A MiniBooNE
confirmation of LSND would
be another revolution.
Too good to be true?**

System

Atmospheric

LSND

Channel

$\rightarrow \nu_\mu$

$\mu \rightarrow \bar{\nu}_e$

$\delta m^2 / \text{eV}^2$

1.2-2 or 6.5

Status

Confirmed

Test

2004 ?

Interpretation

Evidence

neutrino
T violation ... ?


Simplest
interpre-
tation

Three eigenstates with
 $m_1 < m_2 \ll m_3 \sim$
 $m_1 \sim m_2 \sim m_3 \gg$

with
(hierarchical)
 ν (degenerate)


Experimental or
statistical
fluke

Kamland Reactor Neutrino Experiment (Japan)




Japanese nuclear reactors
60 GW (20% world capacity)

- Without Oscillations
2 Neutrino captures / day
- Data taking since
22 January 2002




Kamland Reactor Neutrino Experiment (Japan)



Japanese nuclear reactors
60 GW (20% world capacity)

- Without Oscillations
2 Neutrino captures / day
- Data taking since
22 January 2002



First Kamland results earlier this week
at "Texas in Tuscany" and hep-ex/0212021

Three-Flavor Neutrino Parameters (Ignoring LSND)

Atmospheric
 $32^\circ < \theta_{23} < 60^\circ$

Chooz Limit
 $\theta_{13} < 14^\circ$


Solar
 $27^\circ < \theta_{12} < 41^\circ$

3σ ranges
 hep-ph/0211054

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} 1 & & \\ & c_{23} & s_{23} \\ & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & e^{-i\delta} s_{13} & \\ & 1 & \\ -e^{i\delta} s_{13} & & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} \\ -s_{12} & c_{12} \\ & & 1 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

$c_{12} = \cos \theta_{12}$ etc., δ CP-violating phase

Solar
 24 - 240
 Atmospheric
 1400 - 6000
 $\Delta m^2 / \text{meV}^2$




Cosmology

Tasks and Open Questions

- Precision for θ_{12} and θ_{23} ($\theta_{12} < 45^\circ$ and $\theta_{23} = 45^\circ$?)
- How large is θ_{13} ?
- CP-violating phase?
- Mass ordering? (normal vs inverted)
- Absolute masses? (hierarchical vs degenerate)
- Dirac or Majorana?

Tritium Endpoint Spectrum



Tritium Beta Decay



Endpoint energy 18.6 keV

Mainz Experiment, PLB 460 (1999) 219

$m < 2.8 \text{ eV}$ (95% CL)

Troitsk Experiment, ibid. 227

$m < 2.5 \text{ eV}$ (95% CL)

These
experiments
have reached

2.2 eV
(Neutrino 2002)

Scaled-up spectrometer (KATRIN) may reach 0.3 eV
Currently in preparation - Results in > 5 years

Cosmological Limit on Neutrino Masses

Cosmic neutrino "sea" $\sim 112 \text{ cm}^{-3}$ neutrinos + anti-neutrinos per flavor

$$\Omega_\nu h^2 = \sum \frac{m_\nu}{94 \text{ eV}} < 0.4$$

$$m_\nu < 40 \text{ eV}$$

For all
stable flavors

REST MASS OF MUONIC NEUTRINO AND COSMOLOGY

S. S. Gershtein and Ya. B. Zel'dovich

Submitted 4 June 1966

ZhETF Pis'ma 4, No. 5, 174-177, 1 September 1966


A classic paper:
Gershtein & Zeldovich
JETP Lett. 4 (1966) 120

Low-accuracy experimental estimates of the rest mass of the neutrino [1] yield $m(\nu_e) < 200 \text{ eV}/c^2$ for the electronic neutrino and $m(\nu_\mu) < 2.5 \times 10^6 \text{ eV}/c^2$ for the muonic neutrino.

Cosmological considerations connected with the hot model of the Universe [2] make it possible to strengthen greatly the second inequality. Just as in the paper by Ya. B. Zel'dovich and Ya. A. Smorodinskii [3], let us consider the gravitational effect of the neutrinos on the dynamics of the expanding Universe. The age of the known astronomical objects is not smaller than 5×10^9 years, and Hubble's constant H is not smaller than 75 km/sec-Mpc = $(13 \times 10^9 \text{ years})^{-1}$. It follows therefore that the density of all types of matter in the Universe is at the present time ¹⁾

$$\rho < 2 \times 10^{-28} \text{ g/cm}^3.$$

Neutrino Mass Limit from 2dF Galaxy Survey



Elgaroy et al.,
astro-ph/0204152

$\Sigma m_\nu < 2.2 \text{ eV}$
at 95% CL
(statistical)


Depends on priors for
other parameters

Similar limits based
on the 2dF survey

- Hannestad
astro-ph/0205223
- Lewis & Bridle
astro-ph/0205436

PLANCK Sensitivity to Neutrino Mass

Maximum **polarization multipole** that can be measured to cosmic-variance precision



1 σ sensitivity to Σm_ν

0.07 eV

0.13 eV



0.25 eV

Maximum **temperature multipole** that can be measured to cosmic-variance precision

Steen Hannestad
astro-ph/0211106

Mass-Energy-Inventory of the Universe

Assuming $h = 0.75$




How Many Relic Neutrinos?

Standard thermal population in one flavor $n_{\nu\bar{\nu}} = \frac{3}{11}n_\gamma \approx 112 \text{ cm}^{-3}$

Additional active neutrinos beyond standard population of ν_e, ν_μ, ν_τ	Additional families	Excluded by Z^0 width ($N_\nu = 3$)
	Chemical potentials for ν_e, ν_μ, ν_τ	Possible
Sterile (right-handed) states Populated by $\nu_L \rightarrow \nu_R$ transitions	Dirac mass	Not effective in eV range
	Right-handed currents	Excluded by energy loss of SN 1987A
	Electromagnetic dipole moments	Excluded by energy loss of globular cluster stars
	Oscillations/collisions	Hot/warm/cold DM possible

BBN Limits on Neutrino Flavors



- At BBN one flavor contributes about 16% to cosmic mass-energy density
- Extra flavors modify expansion parameter accordingly

Conservative limit

$$|\Delta N_{\text{eff}}| < 1$$

Burles, Nollett & Turner, astro-ph/9903300

BBN and Neutrino Chemical Potentials

Expansion Rate
Effect
(all flavors)

Energy density in one neutrino flavor with
degeneracy parameter $\xi = \eta/T$

$$\rho_{\nu\bar{\nu}} = \frac{7\pi^2}{120} T_\nu^4 \left[1 + \underbrace{\frac{30}{7} \left(\frac{\xi}{\pi} \right)^2 + \frac{15}{7} \left(\frac{\xi}{\pi} \right)^4}_{\Delta N_{\text{eff}}} \right]$$

Beta equilibrium
effect for
electron flavor




Helium abundance essentially fixed by
 n/p ratio at beta freeze-out

$$\frac{n}{p} = e^{-(m_n - m_p)/T - \xi_{\nu_e}}$$

Effect on helium equivalent to $\Delta N_{\text{eff}} \sim -18 \xi_{\nu_e}$

$$|\Delta N_{\text{eff}}| < 1$$



$$|\xi_{\nu_e}| < 0.06$$

- ν_e beta effect can compensate expansion-rate effect of $\nu_{\mu,\tau}$
- No significant BBN limit on neutrino number density

Chemical Potentials and Flavor Oscillations



Flavor mixing
(neutrino oscillations)

Flavor lepton numbers
not conserved

Only one common nu
chemical potential

Stringent ξ_{ν_e} limit
applies to all flavors
 $|\xi_{\nu_e, \mu, \tau}| < 0.07$

Extra neutrino density
 $\Delta N_{\text{eff}} < 0.0064$

Cosmic neutrino density
close to standard value

Flavor equilibrium before n/p
freeze out?

yes

Solar LMA solution

maybe

LOW (depends on Θ_{13})

no

SMA or VAC

- Our knowledge of the cosmic nu density depends on the solution of the solar neutrino problem
- KamLAND most relevant experiment

- Lunardini & Smirnov, hep-ph/0012056
- Dolgov, Hansen, Pastor, Petcov, Raffelt & Semikoz, hep-ph/0201287
- Abazajian, Beacom & Bell, astro-ph/0203442
- Wong, hep-ph/0203180

Baryogenesis in the Early Universe



Andrei Sakharov
1921 - 1989

Sakharov conditions for creating the
Baryon **A**symmetry of the **U**niverse (**BAU**)

- **C** and **CP** violation
- **Baryon number violation**
- **Deviation from thermal equilibrium**

Particle-physics standard model

- **Violates C and CP**
- **Violates B and L by EW instanton effects**
(**B - L conserved**)

- However, electroweak baryogenesis not quantitatively possible within particle-physics standard model
- Works in SUSY models for small range of parameters

A.Riotto & M.Trodden: Recent progress in baryogenesis
Ann. Rev. Nucl. Part. Sci. 49 (1999) 35

Leptogenesis by Majorana Neutrino Decays

A classic paper

Volume 174, number 1

PHYSICS LETTERS B

26 June 1986

BARYOGENESIS WITHOUT GRAND UNIFICATION

M. FUKUGITA

Research Institute for Fundamental Physics, Kyoto University, Kyoto 606, Japan

and

T. YANAGIDA

*Institute of Physics, College of General Education, Tohoku University, Sendai 980, Japan
and Deutsches Elektronen-Synchrotron DESY, D-2000 Hamburg, Fed. Rep. Germany*

Received 8 March 1986

A mechanism is pointed out to generate cosmological baryon number excess without resorting to grand unified theories. The lepton number excess originating from Majorana mass terms may transform into the baryon number excess through the unsuppressed baryon number violation of electroweak processes at high temperatures.


See-Saw Model for Neutrino Masses

Dirac masses
from coupling
to standard
Higgs field ϕ

Charged Leptons



Neutrinos



Heavy
Majorana
masses
 $M_j > 10^{10} \text{ GeV}$

Lagrangian for
particle masses

$$\mathcal{L}_{\text{mass}} = -\bar{\ell}_L \phi g_\ell e_R - \bar{\ell}_L \phi g_\nu N_R - \frac{1}{2} \overline{N_R^c} M N_R + \text{h.c.}$$


Light Majorana mass

$$(\bar{\nu}_L \quad \bar{N}_R) \begin{pmatrix} 0 & g_\nu \langle \phi \rangle \\ g_\nu \langle \phi \rangle & M \end{pmatrix} \begin{pmatrix} \nu_L \\ N_R \end{pmatrix}$$

Diagonalize

$$(\bar{\nu}_L \quad \bar{N}_R) \begin{pmatrix} \frac{g_\nu^2 \langle \phi \rangle^2}{M} & 0 \\ 0 & M \end{pmatrix} \begin{pmatrix} \nu_L \\ N_R \end{pmatrix}$$

Leptogenesis by Out-of-Equilibrium Decay



CP-violating decays by interference of tree-level with one-loop diagram



$$\Gamma_{\text{Decay}} = g_v^2 \frac{M}{8\pi}$$

W. Buchmüller & M. Plümacher: Neutrino masses and the baryon asymmetry
Int. J. Mod. Phys. A15 (2000) 5047-5086

Leptogenesis by Majorana Neutrino Decays

In see-saw models for neutrino masses, out-of-equilibrium decay of right-handed heavy Majorana neutrinos provides source for CP- and L-violation

Cosmological evolution:

- $B = L = 0$ early on
- Thermal freeze-out of heavy Majorana neutrinos
- Out-of-equilibrium CP-violating decay creates net L
- Shift L excess into B by sphaleron effects

Sufficient deviation from equilibrium distribution of heavy Majorana neutrinos at freeze-out



Limits on Yukawa couplings



Limits on masses of ordinary neutrinos

Requires Majorana neutrino masses below 0.2 eV

Buchmüller, Di Bari & Plümacher, PLB 547 (2002) 128 [hep-ph/0209301]

Leptogenesis - A Popular Research Topic

Fukugita & Yanagida
PLB 174 (1986) 45

Langacker, Peccei & Yanagida
Mod. Phys. Lett. A 1 (1986) 541

Campbell, Davidson & Olive
NPB 399 (1993) 111

Gherghetta & Jungmann
PRD 48 (1993) 1546

Muryama & Yanagida
PLB 322 (1994) 349

Worah
PRD 53 (1996) 3902

Jeannerot
PRL 77 (1996) 3292

Dine, Randall & Thomas
NPB 458 (1996) 291

Buchmüller & Plümacher
PLB 389 (1996) 73

Lazarides, Schaefer & Shafi
PRD 56 (1997) 1324

Ma & Sarkar
PRL 80 (1998) 5716

Plümacher
NPB 530 (1998) 207

Flanz & Paschos
PRD 58 (1998) 113009

Lazarides & Shafi
PRD 58 (1998) 071702

Akhmedov, Rubakov & Smirnov
PRL 81 (1998) 1562

Carlier, Frère & Ling
PRD 60 (1999) 096003

Giudice, Peloso, Riotto & Tkachev
JHEP 9908 (1999) 014

Berger & Brahmachari
PRD 60 (1999) 073009

Ellis, Lola & Nanopoulos
PLB 452 (1999) 87

Barbieri, Creminelli, Strumia & Tetradis
NPB 575 (2000) 61

Frère, Ling, Tytgat & v.Elewyck
PRD 60 (1999) 016005

Dick, Lindner, Ratz & Wright
PRL 84 (2000) 4039

Lalakulich, Paschos & Flanz
PRD 62 (2000) 053006

Asaka, Hamaguchi, Kawasaki & Yanagida
PRD 61 (2000) 083512

Berger
PRD 62 (2000) 013007

Hambye, Ma & Sarkar
PRD 62 (2000) 015010

Mangano & Miele
PRD 62 (2000) 063514

Goldberg
PLB 474 (2000) 389

Rangarajan & Mishra
PRD 61 (2000) 043509

Hirsch & King
PRD 64 (2001) 113005

Falcone & Tramontano
PRD 63 (2001) 073007

Bastero-Gil & King
PRD 63 (2001) 123509

Joshiyura, Paschos & Rodejohann
NPB 611 (2001) 227


Branco, Morozumi, Nobre & Rebelo
NPB 617 (2001) 475

Hambye, Ma & Sarkar
NPB 602 (2001) 23


AND MANY MORE ...

Neutrinoless $\beta\beta$ Decay


0ν mode, enabled by Majorana mass



Standard 2ν mode



Some nuclei decay only by the $\beta\beta$ mode, e.g.




Half life $\sim 10^{21}$ yr

Measured quantity

$$|m_{ee}| = \left| \sum_{i=1}^N \lambda_i |U_{ei}|^2 m_i \right|$$

Best limit from ^{76}Ge

$$|m_{ee}| < 0.35 \text{ eV}$$




Summary of Current Neutrinoless $\beta\beta$ Decay Limits

$0\nu2\beta$ Experimental Situation

2 main experimental approaches:

- Active Source
- Passive Source



Best $0\nu2\beta$ results involve active source experiments

Experiment	Isotope	$T_{1/2}^{0\nu}$ (y)	$\langle m_\nu \rangle$ (eV)
You Ke et al. 1998	^{48}Ca	$> 9.5 \times 10^{21}$ (76%)	< 8.3
Klapdor-Kleingrothaus 2001	^{76}Ge	$> 1.9 \times 10^{25}$	< 0.35
Aalseth et al 2002		$> 1.57 \times 10^{25}$	$< 0.33 - 1.35$
Elliott et al. 1992	^{82}Se	$> 2.7 \times 10^{22}$ (68%)	< 5
Ejiri et al. 2001	^{100}Mo	$> 5.5 \times 10^{22}$	< 2.1
Danevich et al. 2000	^{116}Cd	$> 7 \times 10^{22}$	< 2.6
Bernatowicz et al. 1993	$^{130/128}\text{Te}^*$	$(3.52 \pm 0.11) \times 10^{-4}$	$< 1.1 - 1.5$
Bernatowicz et al. 1993	$^{128}\text{Te}^*$	$> 7.7 \times 10^{24}$	$< 1.1 - 1.5$
Mi DBD – ν 2002	^{130}Te	$> 2.1 \times 10^{23}$	$< 0.85 - 2.1$
Luescher et al. 1998	^{136}Xe	$> 4.4 \times 10^{23}$	$< 1.8 - 5.2$
Belli et al. 2001	^{136}Xe	$> 7 \times 10^{23}$	$< 1.4 - 4.1$
De Silva et al. 1997	^{150}Nd	$> 1.2 \times 10^{21}$	< 3
Danevich et al. 2001	^{160}Gd	$> 1.3 \times 10^{21}$	< 26

Summary of Future $\beta\beta$ Decay Projects

O.Cremonesi at Neutrino 2002


Future projects

Experiment	Author	Isotope	Detector description	$T_{1/2}^{5y}$ (y)	$\langle m_\nu \rangle^*$
COBRA	Zuber 2001	^{130}Te	10 kg CdTe semiconductors	1×10^{24}	0.71
CUORICINO	Arnaboldi et al 2001	^{130}Te	40 kg of TeO_2 bolometers	1.5×10^{25}	0.19
NEMO3	Sarazin et al 2000	^{100}Mo	10 kg of bb(0n) isotopes (7 kg Mo) with tracking	4×10^{24}	0.56
CUORE	Arnaboldi et al. 2001	^{130}Te	760 kg of TeO_2 bolometers	7×10^{26}	0.027
EXO	Danevich et al 2000	^{136}Xe	1 t enriched Xe TPC	8×10^{26}	0.052
GEM	Zdesenko et al 2001	^{76}Ge	1 t enriched Ge diodes in liquid nitrogen + water shield	7×10^{27}	0.018
GENIUS	Klapdor-Kleingrothaus et al 2001	^{76}Ge	1 t enriched Ge diodes in liquid nitrogen	1×10^{28}	0.015
MAJORANA	Aalseth et al 2002	^{76}Ge	0.5 t enriched Ge segmented diodes	4×10^{27}	0.025
DCBA	Ishihara et al 2000	^{150}Nd	20 kg enriched Nd layers with tracking	2×10^{25}	0.035
CAMEO	Bellini et al 2001	^{116}Cd	1 t CdWO_4 crystals in liquid scintillator	$> 10^{26}$	0.069
CANDLES	Kishimoto et al	^{48}Ca	several tons of CaF_2 crystal in liquid scintillator	1×10^{26}	
GSO	Danevich 2001	^{160}Gd	2 t $\text{Gd}_2\text{SiO}_5:\text{Ce}$ cristal scintillator in liquid scintillator	2×10^{26}	0.065
MOON	Ejiri et al 2000	^{100}Mo	34 t natural Mo sheets between plastic scintillator	1×10^{27}	0.036
Xe	Caccianiga et al 2001	^{136}Xe	1.56 t of enriched Xe in liquid scintillator	5×10^{26}	0.066
XMASS	Moriyama et al 2001	^{136}Xe	10 t of liquid Xe	3×10^{26}	0.086

* Staudt, Muto, Klapdor-Kleingrothaus Europh. Lett 13 (1990) 31

Matter Inventory of the Universe

Dark Energy ~ 70%
(Cosmological Constant?)



Baryonic Matter ~ 5%
(~10% of this luminous)

Dark Matter
~ 25%

Neutrinos
min. 0.1%
max. 6%